

GUIDANCE REFERENCES

1: Buising KL, Thursky KA, Black JF, MacGregor L, Street AC, Kennedy MP, Brown GV. A prospective comparison of severity scores for identifying patients with severe community acquired pneumonia: reconsidering what is meant by severe pneumonia. *Thorax*. 2006 May;61(5):419-24. Epub 2006 Jan 31.

2: Thursky KA, Buising KL, Bak N, Macgregor L, Street AC, Macintyre CR, Presneill JJ, Cade JF, Brown GV. Reduction of broad-spectrum antibiotic use with computerized decision support in an intensive care unit. *Int J Qual Health Care*. 2006 Jun;18(3):224-31. Epub 2006 Jan 13.

3: Leong CL, Buising K, Richards M, Robertson M, Street A. Providing guidelines and education is not enough: an audit of gentamicin use at The Royal Melbourne Hospital. *Intern Med J*. 2006 Jan;36(1):37-42.

4: Buising KL, Thursky KA, Bak N, Skull S, Street A, Presneill JJ, Cades JF, Brown GV. Antibiotic prescribing in response to bacterial isolates in the intensive care unit. *Anaesth Intensive Care*. 2005 Oct;33(5):571-7.

5: Buising KL, Thursky KA, Black JF, Brown GV. Are the Australian guidelines asking too much of the Pneumonia Severity Index (PSI)? *Med J Aust*. 2004 May 3;180(9):486-7.

6: Richards MJ, Robertson MB, Dartnell JG, Duarte MM, Jones NR, Kerr DA, Lim LL, Ritchie PD, Stanton GJ, Taylor SE. Impact of a web-based antimicrobial approval system on broad-spectrum cephalosporin use at a teaching hospital. *Med J Aust*. 2003 Apr 21;178(8):386-90.